

Phospho-Rb (Ser807/811) (D9) rabbit mAb

www.abwizbio.com

Support: info@abwizbio.com

Order: sales@abwizbio.com


Catalog: #2051

Store at: -20°C


For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry, WB	Anti-Rabbit IgG	Monoclonal	Rabbit IgGk


Format:	Unconjugated
Cross Reactivity:	Predicted to work with mouse, rat and other homologues.
Formulation:	1X PBS, 0.02% NaN ₃ , 50% Glycerol, 0.1% BSA
Preparation:	Protein A+G
Reactivity:	Human, Mouse
Recommended Usage:	Use 1 ug/mL for flow cytometry and 0.1 ug/mL for western blotting.
Immunogen:	A synthetic phospho-peptide corresponding to residues surrounding Ser807/Ser811 of human phospho Rb
Description:	Retinoblastoma protein (Rb, phospho Rb) is a tumor suppressor protein that is inactivated in a number of diverse cancers. The antiproliferative activity of Rb is mediated by its ability to inhibit the transcription of genes that are required for cell cycle progression. Rb contains conserved sites that are phosphorylated by cyclin-dependent kinases (CDKs). CDK phosphorylation typically promotes protein-protein interactions through creation of a phospho-epitope that becomes structured upon binding its target. However Rb phosphorylation disrupts interactions with its binding partners. When it is phosphorylated, phospho Rb is inactivated and allows excessive cell growth that is seen in cancer cells. Sixteen potential sites for CDK-mediated phosphorylation exist in Rb and twelve of these sites have been shown to be phosphorylated in vivo.
References:	<ol style="list-style-type: none">1. Burkhardt DL et al., (2008) Nat Rev Cancer. 8:671-682.2. Knudsen ES (2008) Nat Rev Cancer. 8:714-724.3. Mittnacht S. (1998) Curr Opin Genet Dev. 8:21-27.4. Weinberg RA. (1995) 81:323-330.5. Henley SA et al., (2012) Cell Div., 7:10.



Flow cytometric analysis of U937 cells untreated and unstained as negative control (blue) or untreated and stained (green) or treated with lambda phosphatase and stained (red) using Phospho-Rb (Ser807/811) antibody RBS807S811-D9. at 1 ug/mL. Cat. #2051.



Western blot analysis of NIH3T3 cell extract untreated or treated with 20% FBS using Phospho-Rb (Ser807/Ser811) antibody RbS807S811-D9 at 0.1 µg/mL. Cat. #2051.



Western blot analysis of 3T3 cell extract untreated or treated with 20% FBS using 0.01 $\mu\text{g/mL}$ Phospho-Rb (Ser807/811) antibody RbS807S811-D9 Cat. #2051 or Company C antibody at 0.1 $\mu\text{g/mL}$ (manufacturer's recommended concentration) developed using the same exposure.