## Phospho-PI3 Kinase p85 (Tyr458)/p55 (Tyr199) (1A11) rabbit mAb FITC conjugate


www.abwizbio.com Support: info@abwizbio.com Order: sales@abwizbio.com

Catalog: #2068 Store at: 2-8°C

For Research Use Only. Not For Use In Diagnostic Procedures.

<b>Applications</b> Flow Cytometry	<b>Detection</b> N/A	<b>Clonality</b> Monoclonal	<b>lsotype</b> Rabbit IgGk	
riow Cytometry	NA	Monocional	Nabbit 190k	
Format:	FITC			
Cross Reactivity:	Predicted to work with mouse,	dicted to work with mouse, rat, and other homologues.  PBS, 0.09% NaN3, 0.2% BSA		
Formulation:	1X PBS, 0.09% NaN3, 0.2% BS	4		
Preparation:	Protein A+G			
Reactivity:	Human,Mouse			
Recommended Usage:	For flow cytometric staining, the suggested use of this reagent is 5 $\mu$ L per million cells or 5 $\mu$ L per 100 $\mu$ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.			
Immunogen:	A synthetic phospho-peptide corresponding to residues surrounding Tyr458 of human phospho PI3K p85			
Description:	phosphorylation at the D-3 post activate signaling pathways in kinase B (PKB) is a major down have different specificities and forms. The PH domains of down secondary messengers, causin kinase activation. The PI3K/Akt	phoinositidine 3-kinase (PI3K) targets phosphoinositide lipids for phorylation at the D-3 position to serve as a second messenger molecule to rate signaling pathways in response to extracellular stimuli. Akt/protein se B (PKB) is a major downstream target of PI3K. The multiple forms of PI3K different specificities and different affinities for various phosphatidylinositol s. The PH domains of downstream proteins in these pathways bind to these indary messengers, causing altered cellular distribution and subsequent se activation. The PI3K/Akt pathway is altered in a large proportion of human ers, as this pathway regulates cellular survival, cell cycle progression, and growth.		
References:	Franke TF, Kaplan DR, and Can Fresno JAV, Casado E, Cejas P, 193-204.			





## Phospho-PI3K p85(Y458)/p55(Y199)

Flow cytometric analysis of Ramos cells stained untreated (red) or treated with pervanadate (green) using phospho-PI3 Kinase p85 (Tyr458)/p55 (Tyr199) antibody PI3KY458-1A11 FITC conjugate. Cat. #2068.