Phospho-CrkL (Tyr207) (G4) rabbit mAb SureLight488 Conjugate

www.abwizbio.com

Support: info@abwizbio.com Order: sales@abwizbio.com

#2095

Store at: 2-8°C

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry	y N/A	Monoclonal	Rabbit IgGk
Format:	SureLight 488		
Cross Reactivity:	Predicted to work with mouse, rat and other homologues.		
Formulation:	1X PBS, 0.09% NaN3, 0.2% BSA		
Preparation:	Protein A+G		
Reactivity:	Human,Mouse		
Recommended Usage:	For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells or 5 μ L per 100 μ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.		
Immunogen:	A synthetic phospho0peptide corresponding to residues surrounding Tyr207 of human phospho Crkl		
Description:	CrkL (v-Crk sarcoma virus CT10 oncogene-like protein) is an adaptor protein composed of one Src Homology 2 (SH2) and two Src Homology 3 (SH3) domains separated by flexible linker sequences that act as building blocks to assemble multiprotein complexes (1). The Crk adaptor proteins (Crk and CrkL) constitute an integral part of a network of essential signal transduction pathways in humans and other organisms that act as major convergence points in tyrosine kinase signaling. CRKL is required for the normal development of multiple tissues that rely on fibroblast growth factor 8 (FGF8). Phosphorylation of Crk on Tyr 221 or CrkL on Tyr 207 causes intramolecular binding of the linker region to the SH2 domain, sequestering the SH2 and SH3N and preventing them from binding target proteins (2,3). Mounting evidence indicates that dysregulation of Crk proteins is associated with human diseases, including cancer and susceptibility to pathogen infections.		
References:	 ten Hoeve, J., C. Morris, N. Heisterkamp, and J. Groffen. 1993. Isolation and chromosomal localization of CRKL, a human Crk-like gene. Oncogene 82469-2474. Rosen MK, Yamazaki T, Gish GD, Kay CM, Pawson T, Kay LE: Direct demonstration of an intramolecular SH2-phosphotyrosine interaction in the Crk protein. Nature. 1995, 374: 477-479. Kobashigawa Y, Sakai M, Naito M, Yokochi M, Kumeta H, Makino Y, Ogura K, Tanaka S, Inagaki F: Structural basis for the transforming activity of human cancer-related signaling adaptor protein CRK. Nat Struct Mol Biol. 2007, 14: 503-510. 		





Events

Flow cytometric analysis of K562 human myelogenous leukemia cells, treated with imatinib and unstained as negative control (blue) or treated with imatinib (red) or treated with pervanadate (green) and stained using Phospho-CRKL (Tyr207) antibody SureLight488 conjugate, CRKLY207-G4. Cat. #2095.

Phospho-CRKL (Y207) SL



©2018 Abwiz Bio 9823 Pacific Heights Blvd., Suite J, San Diego, CA 92121