

Phospho-KSR1 (Ser392) (3A4) rabbit mAb APC conjugate

www.abwizbio.com

Support: info@abwizbio.com

Order: sales@abwizbio.com


Catalog: #2189

Store at: 2-8°C

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry	N/A	Monoclonal	Rabbit IgGκ

Format:	APC
Cross Reactivity:	Predicted to work with mouse, rat and other homologues.
Formulation:	1X PBS, 0.09% NaN ₃ , 0.2% BSA
Preparation:	Protein A+G
Reactivity:	Human
Recommended Usage:	For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells or 5 µL per 100 µL of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.
Immunogen:	A synthetic phosphor-peptide corresponding to residues surrounding Ser392 of human phospho KSR1
Description:	
References:	1. Kortum RL, et al., (2004) Mol. Cell. Biol. 24:4407-16. 2. Nguyen A, et al., (2002) Mol. Cell. Biol. 22:3035-45. 3. Roy F, et al., (2002) Genes Dev. 16:427-38. 4. Kortum RL, (2006) Cell. Biol. 26:2202-14. 5. Lozano J, et al., (2003) Cancer Res. 63:4232-8. 6. Sundaram M, et al., (1995) Cell 83:889-901. 7. McKay MM, et al., (2009) Proc. Natl. Acad. Sci. U. S. A. 106:11022-7. 8. Therrien, M, et al., (1995) Cell 83, 879-88.



Flow cytometric analysis of Hela cells untreated (green) or treated with phosphatase (red) using Phospho-KSR1(Ser392) (3A4) Rabbit mAb (APC Conjugate) KSP1S392-3A4 #2189, or concentration-matched Rabbit (G9) mAb IgG Isotype Control (APC Conjugate) #2144 for cells untreated (black) or treated with phosphatase (blue).