

Phospho-PLC γ 1 (Tyr783) (C4) rabbit mAb

www.abwizbio.com

Support: info@abwizbio.com

Order: sales@abwizbio.com


Catalog: #2201

Store at: -20°C






For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry	Anti-Rabbit IgG	Monoclonal	Rabbit IgGk


Format:	Unconjugated
Cross Reactivity:	Predicted to work with mouse, rat, and other homologues.
Formulation:	1X PBS, 0.02% NaN ₃ , 50% Glycerol, 0.1% BSA
Preparation:	Protein A+G
Reactivity:	Human, Mouse
Recommended Usage:	1 μ g/mL - 0.001 μ g/mL. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.
Immunogen:	A synthetic phospho-peptide corresponding to residues surrounding Tyr783 of human phospho PLC γ 1.
Description:	<p>The Phospholipase C (PLC) isozymes hydrolyze phosphatidyl inositolphosphate to inositol triphosphate and diacylglycerol. In response to extracellular stimuli such as hormones, growth factors and neurotransmitters, PLC hydrolyzes phosphatidylinositol 4,5-bisphosphate (PIP₂) to generate diacylglycerols (DAGs) and water-soluble phosphorylated derivatives, such as inositol 1,4,5-triphosphate (IP₃). Within the PLC family, PLCγ is the only member that contains SH2 and SH3 domains, necessary for phospho PLCγ activation. Phospho PLCγ, upon activation, can interact with receptor tyrosine kinases.</p>
References:	<ol style="list-style-type: none">1. Singer, W.D. et al. (1997) Annu. Rev. Biochem. 66, 475-509.2. Hernandez D, et al. (1994) Genomics 23 (2): 504-507.3. Smrcka, A.V. et al. (1991) Science 251, 804-807.4. Taylor, S.J. et al. (1991) Nature 350, 516-518.




Phospho-PLCG1(Y783)

	SWELLID	Treatment	Median : BL1 - A
	C4+PP	PV	1188
	C4+NP	PV	27480
	C4 0.05 ug/mL	PV	24493
	C4 0.05 ug/mL	imatinib	1784
	2'Ab	imatinib	699

Peptide blockage flow cytometric analysis of Hela cells secondary antibody only negative control (blue) treated with imatinib (red) treated with pervanadate (green) treated with PV + blocked with non-phospho-peptide (violet) or treated with PV + blocked with phospho-peptide (brown) using Phospho-PLCγ1 (Tyr783) antibody at 0.05 µg/mL PLCγ1Y783-C4. Cat. #2201.




PLCg1Y783-C4 recognizes basal phosphorylation levels in mouse cells. Flow cytometric analysis of L929 cells secondary antibody only (blue) or 0.1 $\mu\text{g/mL}$ of isotype control Cat. #2141 (orange) or of Phospho-PLC γ 1 (Tyr783) antibody PLCg1Y783-C4 (green) Cat. #2201.



Flow cytometric analysis of HeLa cells secondary antibody only negative control (blue) or treated with imatinib (red) or with pervanadate (green) using 0.01 $\mu\text{g/mL}$ Phospho-PLC γ 1 (Tyr783) antibody PLCg1Y783-C4. Cat. #2201.

Abwiz Cat. #2201


0.1 µg/mL



Phospho-PLCγ1 (Y783)

Company C

0.1 µg/mL



Phospho-PLCγ1 (Y783)

Flow cytometric analysis of HeLa cells secondary antibody only negative control (blue) or treated with imatinib (red) or with pervanadate (green) using Phospho-PLCγ1 (Tyr783) antibody PLCγ1Y783-C4 (Abwiz Cat. #2201) or Company C antibody at 0.1 ug/mL (manufacturer's recommended concentration).