

# Phospho-PLC $\beta$ 1 (Tyr783) (C4) rabbit mAb

www.abwizbio.com

Support: info@abwizbio.com

Order: sales@abwizbio.com

## #2201

Store at: -20°C

**For Research Use Only. Not For Use In Diagnostic Procedures.**

Applications	Detection	Clonality	Isotype
Flow Cytometry	Anti-Rabbit IgG	Monoclonal	Rabbit IgGk

**Format:** Unconjugated

**Cross Reactivity:** Predicted to work with mouse, rat, and other homologues.

**Formulation:** 1X PBS, 0.02% NaN<sub>3</sub>, 50% Glycerol, 0.1% BSA

**Preparation:** Protein A+G

**Reactivity:** Human, Mouse

### Recommended


**Usage:** 1 $\mu$ g/mL ? 0.001 $\mu$ g/mL. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.

**Immunogen:** A synthetic phospho-peptide corresponding to residues surrounding Tyr783 of human phospho PLC $\beta$ 1.

**Description:** The Phospholipase C (PLC) isozymes hydrolyze phosphatidyl inositolphosphate to inositol triphosphate and diacylglycerol. In response to extracellular stimuli such as hormones, growth factors and neurotransmitters, PLC hydrolyzes phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>) to generate diacylglycerols (DAGs) and water-soluble phosphorylated derivatives, such as inositol 1,4,5-triphosphate (IP<sub>3</sub>). Within the PLC family, PLC $\beta$  is the only member that contains SH2 and SH3 domains, necessary for phospho PLC $\beta$  activation. Phospho PLC $\beta$ , upon activation, can interact with receptor tyrosine kinases.

### References:


1. Singer, W.D. et al. (1997) Annu. Rev. Biochem. 66, 475-509.
2. Hernandez D, et al. (1994) Genomics 23 (2): 504-507.
3. Smrcka, A.V. et al. (1991) Science 251, 804-807.
4. Taylor, S.J. et al. (1991) Nature 350, 516-518.



Peptide blockage flow cytometric analysis of HeLa cells secondary antibody only negative control (blue) treated with imatinib (red) treated with pervanadate (green) treated with PV + blocked with non-phospho-peptide (violet) or treated with PV + blocked with phospho-peptide (brown) using Phospho-PLC $\gamma$ 1 (Tyr783) antibody at 0.05  $\mu$ g/mL PLC $\gamma$ 1Y783-C4. Cat. #2201.

PLC $\gamma$ 1Y783-C4 recognizes basal phosphorylation levels in mouse cells. Flow cytometric analysis of L929 cells secondary antibody only (blue) or 0.1  $\mu$ g/mL of isotype control Cat. #2141 (orange) or of Phospho-PLC $\gamma$ 1 (Tyr783) antibody PLC $\gamma$ 1Y783-C4 (green) Cat. #2201.

Flow cytometric analysis of HeLa cells secondary antibody only negative control (blue) or treated with imatinib (red) or with pervanadate (green) using 0.01  $\mu$ g/mL Phospho-PLC $\gamma$ 1 (Tyr783) antibody PLC $\gamma$ 1Y783-C4. Cat. #2201.



Flow cytometric analysis of HeLa cells secondary antibody only negative control (blue) or treated with imatinib (red) or with pervanadate (green) using Phospho-PLC $\gamma$ 1 (Tyr783) antibody PLC $\gamma$ 1Y783-C4 (Abwiz Cat. #2201) or Company C antibody at 0.1  $\mu$ g/mL (manufacturer's recommended concentration).