



For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry	Anti-Rabbit IgG	Monoclonal	Rabbit IgGk

<b>Format:</b>	Unconjugated
<b>Cross Reactivity:</b>	Predicted to work with mouse, rat and other homologues.
<b>Formulation:</b>	1X PBS, 0.02% NaN <sub>3</sub> , 50% Glycerol, 0.1% BSA
<b>Preparation:</b>	Protein A+G
<b>Reactivity:</b>	Human, Mouse
<b>Recommended Usage:</b>	1μg/mL – 0.001μg/mL. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.
<b>Immunogen:</b>	A synthetic phospho-peptide corresponding to residues surrounding Tyr693 of human phospho Stat4
<b>Description:</b>	In response to IL-12 binding, the IL-12 receptor activates the Jak kinases, which phosphorylate tyrosine residues of IL-12R $\beta$ 2. These phosphorylated receptors recruit Stat4 through its SH2 domain, whereupon Stat4 is phosphorylated at Tyr693 in its C-terminal transactivation domain. Phosphorylation promotes Stat4 homodimerization and translocation to the nucleus, where it promotes gene transcription. The N-terminal domain of Stat4 appears to be required for maximal stabilization and for the binding of Stat4 dimers to lower-affinity DNA binding sites. Stat4-deficient mice have demonstrated that this gene is required to both promote Th1 development and inhibit Th2 differentiation due to disabling IL-12 receptor-mediated responses.
<b>References:</b>	Kaplan MH, Sun Y, Hoey T, and Grusby MJ. (1996) Nature. 382: 174-177. Chang H, Zhang S, Oldham I, Naeger L, Hoey T, and Kaplan MH. (2003) Journal of Biological Chemistry. 278: 32471-32477.




Flow cytometric analysis of K562 cells unstained and treated with imatinib as negative control (blue) or treated with imatinib and stained (red) or treated with IFN $\alpha$  + IL-4 + pervanadate and stained (green) using Phospho-Stat4 (Tyr693) antibody Stat4Y693-F6 at 0.01  $\mu$ g/mL. Cat #2281.




Flow cytometric analysis of NIH3T3 cells secondary antibody only negative control (blue) or treated with imatinib (red) or with pervanadate (green) using Phospho-Stat4 (Tyr693) antibody Stat4Y693-F6 at 0.1  $\mu$ g/mL. Cat #2281.


**Abwiz Cat. #2281**  
**0.01  $\mu$ g/mL**



**Company C**  
**0.1  $\mu$ g/mL**  
(recommended conc.)



Flow cytometric analysis of K562 cells secondary antibody only negative control (blue) or treated with imatinib (red) or with pervanadate (green) using 0.01  $\mu$ g/mL of Phospho-Stat4 (Tyr693) antibody Stat4Y693-F6 Cat. #2281 or Company C at 0.1  $\mu$ g/mL (manufacturer's recommended concentration).



	SampleID	Treatment	Peptide Block	Median : BL1.A
■	F6	IFN	Non-phos.	6432
■	F6	Ctrl	Non-phos.	3583
■	F6	IFN	Phospho.	1119
■	F6	Ctrl	Phospho.	665
■	F6	IFN	-	6146
■	F6	Ctrl	-	3481
■	2' only	Ctrl	-	551

Peptide blocking flow cytometric analysis of K562 cells secondary antibody only negative control (light blue) or untreated (grey) or IFN $\alpha$  + IL-4 + pervanadate-treated (orange) using 0.1  $\mu$ g/mL isotype control Cat. #2141 or untreated (red) or treated (green) or untreated and blocked with phospho-peptide (black) or treated and blocked with phospho peptide (gold) or untreated and blocked with non-phospho peptide (dark blue) or treated and blocked with non-phospho peptide (purple) using Phospho-Stat4 (Tyr693) antibody Stat4Y693-F6 at 0.1  $\mu$ g/mL. Cat. #2281.