

# Phospho-BLNK (Tyr84) (H4) rabbit mAb

[www.abwizbio.com](http://www.abwizbio.com)

Support: [info@abwizbio.com](mailto:info@abwizbio.com)

Order: [sales@abwizbio.com](mailto:sales@abwizbio.com)


**Catalog:** #2291

**Store at:** -20°C

*For Research Use Only. Not For Use In Diagnostic Procedures.*


Applications	Detection	Clonality	Isotype
Flow Cytometry	Anti-Rabbit IgG	Monoclonal	Rabbit IgGk

<b>Format:</b>	Unconjugated
<b>Cross Reactivity:</b>	Predicted to work with mouse, rat and other homologues.
<b>Formulation:</b>	1X PBS, 0.02% NaN <sub>3</sub> , 50% Glycerol, 0.1% BSA
<b>Preparation:</b>	Protein A+G
<b>Reactivity:</b>	Human
<b>Recommended Usage:</b>	1µg/mL – 0.001µg/mL. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.
<b>Immunogen:</b>	A synthetic phospho-peptide corresponding to residues surrounding Tyr84 of human phospho BLNK
<b>Description:</b>	BLNK protein, known as SLP-65 play an important role as adaptor protein in B-lineage cells. BLNK associates with proteins in the cytoplasmic side of plasma membrane through its N-terminal leucine zipper motif. Upon BLNK activation on its tyrosine, BLNK binds to Btk, Vav, Brb2, Syk, and HPK1. Through this associations, BLNK mediates Ca <sup>2+</sup> mobilization, for ERK1/2, JNK and p38 MAP kinase activation. After phosphorylation, BLNK binds Btk and PLC <sub>γ</sub> 2 through their SH2 domains and mediates PLC <sub>γ</sub> 2 activation by Btk. BLNK also binds other signaling molecules such as Vav, Grb2, Syk, and HPK1. BLNK plays an important role in BCR-dependent progression of B cell development, BCR-mediated B cell survival, activation, proliferation, and T-independent immune responses.
<b>References:</b>	Fu, C., et al. (1998) Immunity 9: 93–103. Goitsuka, R., et al. (1998) J. Immunol. 161: 5804–5808. Tsuiji, S., et al. (2001) J. Exp. Med. 194: 529–539.



IgG	Treatment	Peptide Block	Median : BL1-A
■ H4	IFNa	Non-phos	88579
■ H4	Ctrl	Non-phos	2748
■ H4	IFNa	Phos	700
■ H4	Ctrl	Phos	343
■ H4	IFNa	-	75009
■ H4	Ctrl	-	4167
■ 2' only	Ctrl	-	352


Peptide blocking flow cytometric analysis of Daudi cells secondary antibody only negative control (light blue) or untreated (red) or with IFNa + IL-4 + pervanadate (green) or untreated and blocked with phospho-peptide (black) or treated and blocked with phospho peptide (gold) or untreated and blocked with non-phospho peptide (dark blue) or treated and blocked with non-phospho peptide (purple) using Phospho-BLNK (Tyr84) antibody BLNKY84-H4 at 0.01 µg/mL. Cat. #2291.



Flow cytometric analysis of Daudi cells secondary antibody only negative control (blue) or untreated (red) or treated with IFNa + IL-4 + pervanadate (green) using Phospho-BLNK (Tyr84) antibody BLNKY84-H4 at 0.01 µg/mL. Cat. #2291.

**Abwiz Cat. #2291**

**0.01 µg/mL**




**Phospho-BLNK (Y84)**

**Company B**

**2.5 µg/mL**

**(recommended conc.)**



**Phospho-BLNK (Y84)**

Flow cytometric analysis of Daudi cells secondary antibody only negative control (blue) or untreated (red) or treated with IFNa + IL-4 + pervanadate (green) using 10 ng/mL of Phospho-BLNK (Tyr84) antibody BLNKY84-H4 (Abwiz Cat. #2291) or Company B antibody at 2.5 ug/mL (manufacturer's recommended concentration).