## Phospho-BLNK (Tyr84) (H4) rabbit mAb SureLight488 conjugate

www.abwizbio.com Support: info@abwizbio.com Order: sales@abwizbio.com


**Store at:** 2-8°C

**Catalog:** #2295

For Research Use Only. Not For Use In Diagnostic Procedures.

<b>Applications</b> Flow Cytometry	Detection N/A	<b>Clonality</b> Monoclonal	<b>lsotype</b> Rabbit IgGk
Format:	SureLight 488		
Cross Reactivity:	Predicted to work with mouse, rat and other homologues.		
Formulation:	1X PBS, 0.09% NaN3, 0.2% BSA		
Preparation:	Protein A+G		
Reactivity:	Human		
Recommended Usage:	For flow cytometric staining, the suggested use of this reagent is 5 $\mu$ L per million cells or 5 $\mu$ L per 100 $\mu$ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.		
Immunogen:	A synthetic phospho-peptide corresponding to residues surrounding Tyr84 of human phospho BLNK		
<b>Description:</b>	BLNK protein, known as SLP-65 play an important role as adaptor protein in B- lineage cells. BLNK associates with proteins in the cytoplasmic side of plasma membrane through its N-terminal leucine zipper motif. Upon BLNK activation on its tyrosine, BLNK binds to Btk, Vav, Brb2, Syk, and HPK1. Through this associations, BLNK mediates Ca2+ mobilization, for ERK1/2, JNK and p38 MAP kinase activation. After phosphorylation, BLNK binds Btk and PLC $\gamma$ 2 through their SH2 domains and mediates PLC $\gamma$ 2 activation by Btk. BLNK also binds other signaling molecules such as Vav, Grb2, Syk, and HPK1. BLNK plays aa important role in BCR-dependent progression of B cell development, BCR-mediated B cell survival, activation, proliferation, and T-independent immune responses.		
References:	Fu, C., et al. (1998) Immunity 9: 93–103. Goitsuka, R., et al. (1998) J. Immunol. 161: 5804–5808. Tsuji, S., et al. (2001) J. Exp. Med. 194: 529–539.		





Phospho-BLNK1 (Y84) SureLight

Flow cytometric analysis of Ramos cells, unstained and untreated cells as negative control (blue) or untreated and stained (red) or treated with INFa+IL-4 +pervanadate and stained (green) using Phospho-BLNK (Tyr84) antibody BLNKY84-H4 SureLight488 conjugate. Cat #2295.

