
Phospho-PDK1 (Ser241) (F7) rabbit mAb APC conjugate

www.abwizbio.com Support: info@abwizbio.com Order: sales@abwizbio.com

Catalog: #2299 Store at: 2-8°C

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications Flow Cytometry	Detection N/A	Clonality Monoclonal	Isotype Rabbit IgGk	
riow Cytometry	N/A	Monocional	Nabbit 190k	
Format:	APC			
Cross Reactivity:	Predicted to work with mouse, rat and other homologues.			
Formulation:	1X PBS, 0.09% NaN3, 0.2% BSA			
Preparation:	Protein A+G	ı+G		
Reactivity:	Human			
Recommended Usage:	For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells or 5 μ L per 100 μ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.			
Immunogen:	A synthetic phospho-peptide c human phospho PDK1	synthetic phospho-peptide corresponding to residues surrounding Ser241 of		
Description:	PDK1 is a Ser/Thr kinase that is ubiquitously expressed throughout human tissues. PDK1 phosphorylates protein kinase B (PKB or Akt) at both Thr308 and Ser473 in vivo. PDK1 is active and phosphorylated in basal conditions, where it exists predominantly in the cytosol with a small fraction at the plasma membrane. This membrane association is likely mediated by PDK1 binding to phospholipids through its PH domain, which has been found to have strong affinity to specific phospholipids. PDK1 is a master regulator of at least 23 related protein kinases, and more than 50% of all human cancers show significant overstimulation of the PDK1 pathway. Most small-molecule inhibitors of PDK1 target the ATP binding site.			
References:	Peifer C and Alessi DR. (2008) Vanhaesebroeck B and Alessi			



Flow cytometric analysis of 293T cells unstained K252a treated cells (blue) or treated with K252a (red) or with pervanadate (green) using phospho-PDK1 (Ser241) antibody PDK1S241-F7 APC conjugate. Cat. #2299. Abwiz Bio ©2026 Abwiz Bio