Phospho-PKCa (Thr497) (F1) rabbit mAb PE Conjugate


www.abwizbio.com Support: info@abwizbio.com Order: sales@abwizbio.com

Catalog: #2337 **Store at:** 2-8^oC

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications Flow Cytometry	Detection N/A	Clonality Monoclonal	Isotype Rabbit IgGk
Format:	PE		
Cross Reactivity:	Predicted to work with mouse, rat and other homologues.		
Formulation:	1X PBS, 0.09% NaN3, 0.2% BSA		
Preparation:	Protein A+G		
Reactivity:	Human,Mouse,Rat		
Recommended Usage:	For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells or 5 μ L per 100 μ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.		
Immunogen:	A synthetic phospho-peptide corresponding to residues surrounding Thr497 of human phospho PKC $\!\alpha$		
Description:	PKC α is a calcium-dependent isozyme of the PKC family that phosphorylates serine/threonine residues in apoptosis and cellular proliferation and differentiation pathways, including the MAPK cascade. PKC α directly phosphorylated Raf-1, inducing survival genes. An increase in PKC α is associated with multi-drug resistance in cancer cell lines, and increased expression in breast cancers is noted as causing a particularly malignant phenotype. Thus PKC α has been the target of novel cancer therapeutics, with some promising developments in microRNA inhibitors. PKC α is itself phosphorylated by mTOR. PKC α also plays an important role in water regulator and solute absorption in the cell, where it regulates aquaporin 2 by initiating AQP2 ubiquination and lysosomal degradation.		
References:	Blobe GC, et al., (1993) JBC. 268:658-664. Sim JH, et al., (2014) PLoS One. 9:e101753. Martin EC, et al. (2012) Molecular Carcinogenesis. 53:38-48.		





Flow cytometric analysis of NIH3T3 cells treated with imatinib and unstained as negative control (blue) or treated with imatinib (red) or treated with pervanadate (green) and stained using PKC α (T497) antibody PKCaT497-F1 PE conjugate. Cat. #2337.