

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry, ELISA	Streptavidin	Monoclonal	Rabbit IgGk

Format: Biotin

Cross Reactivity: Predicted to work with mouse, rat and other homologues.

Formulation: 1X PBS, 0.09% NaN3, 0.2% BSA

Preparation: Protein A+G


Reactivity: Human

Recommended Usage: For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells or 5 µL per 100 µL of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.


Immunogen: A synthetic phospho-peptide corresponding to residues surrounding human Btk (Y551)

Description: Btk is a major node in the B-cell receptor signaling pathway, where it regulates B cell maturation, activation, survival, differentiation, and proliferation. Btk is activated by Src family kinases, including Lyn, which phosphorylates Btk at Tyr551. Upon phosphorylation at this site, Btk is recruited to the plasma membrane where autophosphorylation at Tyr223 occurs. The Btk signaling pathway is a major target of small molecule inhibitors for B-cell lymphoma, autoimmune diseases, and non-Hodgkin's lymphomas. These inhibitors either form a covalent bond at Cys481 in the ATP-binding site or serve as reversible inhibitors that bind the SH3 pocket and stabilize inactive Btk.

References: Liang, C, Tian D, Ren X, et al. (2018) European Journal of Medicinal Chemistry. 151: 315-326.




Flow cytometric analysis of Ramos cells secondary antibody only negative control (blue) or untreated (red) or treated with pervanadate (green) using Phospho-Btk (Tyr551) antibody BtkY551-G12-Biotin conjugate at 0.1 μ g/mL. Cat. #2341B.



Peptide ELISA using BtkY551-G12-Biotin conjugate (Cat. #2341B) titrated starting from 0.3 μ g/mL shows binding to only Btk pY551 phospho peptide and no cross-reactivity to Btk Y551 non-phospho peptide or to control peptides.

BtkY551-G12-Biotin conjugate



Ramos cell lysate, untreated or treated with 1mM Pv for 30 min

Direct ELISA using Ramos cellular lysate coated directly to the plate surface after lysis following no treatment or treatment with 1mM pervanadate for 30 min. ELISA wells were tested in duplicate using 0.12 mg/mL total protein coated lysate and 1 ug/mL BtkY551-G12-Biotin conjugate (Cat. #2341B) IgG.