Phospho-Btk (Tyr551) (G12) rabbit mAb APC conjugate

www.abwizbio.com

Support: info@abwizbio.com Order: sales@abwizbio.com

#2344

Store at: 2-8°C

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry	y N/A	Monoclonal	Rabbit IgGk
Format:	APC		
Cross Reactivity:	Predicted to work with mouse, rat and other	er homologues.	
Formulation:	1X PBS, 0.09% NaN3, 0.2% BSA		
Preparation:	Protein A+G		
Reactivity:	Human		
Recommended Usage:	For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells or 5 μ L per 100 μ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application.		
Immunogen:	A synthetic phospho-peptide correspondin	g to residues surrounding Tyr551	of human phospho Btk
Description:	Btk is a major node in the B-cell receptor signaling pathway, where it regulates B cell maturation, activation, survival, differentiation, and proliferation. Btk is activated by Src family kinases, including Lyn, which phosphorylates Btk at Tyr551. Upon phosphorylation at this site, Btk is recruited to the plasma membrane where autophosphorylation at Tyr223 occurs. The Btk signaling pathway is a major target of small molecule inhibitors for B-cell lymphoma, autoimmune diseases, and non-Hodgkin?s lymphomas. These inhibitors that bind the SH3 pocket and stabilize inactive Btk.		
References:	Liang, C, Tian D, Ren X, et al. (2018) Euro	ppean Journal of Medicinal Chemis	stry. 151: 315-326.



EVENTS

BTK (Y551) APC

Flow cytometric analysis of Ramos cells unstained untreated as negative control (blue) or stained untreated (red) or treated with pervanadate (green) using Phospho-Btk (Tyr551) antibody BtkY551-G12 APC conjugate. Cat. #2344.

