

# Phospho-c-Cbl (Tyr774) (R4C5) rabbit mAb

www.abwizbio.com

Support: info@abwizbio.com

Order: sales@abwizbio.com

## #2361

Store at: -20°C

**For Research Use Only. Not For Use In Diagnostic Procedures.**

Applications	Detection	Clonality	Isotype
Flow Cytometry, WB	Anti-Rabbit IgG	Monoclonal	Rabbit IgGk

**Format:** Unconjugated

**Cross Reactivity:** Predicted to work with mouse, rat and other homologues.

**Formulation:** 1X PBS, 0.02% NaN<sub>3</sub>, 50% Glycerol, 0.1% BSA

**Preparation:** Protein A+G

**Reactivity:** Human, Mouse

### Recommended


**Usage:** 1µg/mL ? 0.001µg/mL. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.

**Immunogen:** A synthetic phospho-peptide corresponding to residues surrounding Tyr774 of human phospho c-Cbl


**Description:** The c-Cbl (Casitas B-lineage Lymphoma) proto-oncogene is a ubiquitously expressed cytoplasmic adaptor protein that contains multiple functional domains, including an amino-terminal tyrosine kinase-binding (TKB) domain, a RING finger motif, and a proline-rich region. The TKB recognizes phosphorylated tyrosines on activated receptor tyrosine kinases (RTKs) and on other nonreceptor tyrosine kinases, while the RING finger motif recruits ubiquitin-conjugating enzymes. These two domains are primarily responsible for the ubiquitin ligase activity of c-Cbl and downregulation of RTKs (1). The proline-rich region contains 14-3-3 protein-binding and SH3 domain-binding motifs. c-Cbl is phosphorylated at Y700, Y731, and Y774 by Syk- and Src-family kinases after the stimulation of some integrins and a wide variety of receptors for immunoglobulins, antigens, hormones, growth factors, and cytokines. Phosphorylated Y774 interacts with the SH2 domain of Crk (1,2). The c-Cbl adapter protein is expressed in the cytoplasm in all tissues, with especially high levels of expression in hematopoietic cells (3,4). Through its many functional sites, c-Cbl plays key roles in the positive and negative regulation of vital cell functions, including T Cell Receptor-mediated cellular immune responses. In human cancer tissues, c-Cbl is frequently tyrosine-phosphorylated in a tumor-specific manner (5).

### References:

1. Christine, B.F. et al. (2001) Nat. Rev. Mol. Cell Biol. 2: 294-307.
2. Feshchenko, E.A. et al. (1998) J. Biol. Chem. 273: 8323-8331.
3. Blake, T.J. et al. (1991) Oncogene. 6: 653-657.
4. Thien, C.B. and Langdon, W.Y. (1998) Immunol. Cell Biol. 76: 473-482.
5. Kamei, T. et al. (2000) Int. J. Oncol. 17: 335-339.




Flow cytometric analysis of Daudi cells secondary antibody only negative control (blue) or untreated (red) or treated with IFN $\gamma$  + IL-4 + pervanadate (green) using Phospho-c-Cbl (Tyr774) antibody CbLY774-R4C5 at 0.01  $\mu$ g/mL. Cat. #2361.



Western blot analysis of extracts from Ramos or Jurkat cells, imatinib or pervanadate treated using Phospho-c-Cbl (Tyr774) antibody c-CbLY774-R4C5 at 0.1  $\mu$ g/mL, Cat#2361


#### **Abwiz Cat. #2361**

**0.1  $\mu$ g/mL**



#### **Company A**

**0.04  $\mu$ g/mL**  
(recommended conc.)



Western blot analysis of Jurkat cell extract treated with imatinib or with pervanadate using 0.1  $\mu$ g/mL Phospho-c-Cbl (Tyr774) antibody CbLY774-R4C5 Cat. #2361 or Company A antibody at 0.04  $\mu$ g/mL (manufacturer's recommended concentration) developed using the same exposure.