

Phospho-SEK1/MKK4 (Ser257) (C5) rabbit mAb PE Conjugate

www.abwizbio.com

Support: info@abwizbio.com

Order: sales@abwizbio.com

#2367

Store at: 2-8°C

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry	N/A	Monoclonal	Rabbit IgGk

Format: PE

Cross Reactivity: Predicted to work with mouse, rat and other homologues.

Formulation: 1X PBS, 0.09% NaN₃, 0.2% BSA

Preparation: Protein A+G

Reactivity: Human, Mouse, Monkey

Recommended


Usage: For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells or 5 µL per 100 µL of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.

Immunogen: A synthetic phospho-peptide corresponding to residues surrounding Ser257 of human phospho SEK1/MKK4

Description: SEK1, in cooperation with MKK7, activates JNK by tyrosine phosphorylation in response to cellular stress signals, including osmotic stress, DNA damage, cytokines, and heat shock (1). SEK1 itself is activated by serine phosphorylation by MAP3Ks, including MEKK1, MLKs, ASK1, and TAK1 (2). In response to TNF α stimulation in melanoma cells, filamin interacts with SEK1 as a scaffold protein (3). SEK1 plays an important role in T cell signaling pathways. Both SEK1-dependent and independent pathways operate for SAPK activation in response to different cellular stresses and stimuli (4). SEK1 also plays an important role in embryonic development. SEK1 may activate c-Jun and c-Fos during mammalian embryonic liver development.

References:

1. Tibbles LA, et al., (1996) The EMBO Journal. 15:7026-7035.
2. Nishina H, et al., (1997) Nature. 385:350-353.
3. Hu L, et al., (2019) The JNK Pathway in Drug Resistance. Targeting Cell Survival Pathways to Enhance Response to Chemotherapy. 87-100.
4. Nishina H, et al. (1999) Development. 126:505-516.



Flow cytometric analysis of 293T cells treated with lambda phosphatase (red) or treated with UV (100 mJ/cm²)+TPA (200nM, 1 hr recovery ; green) using Phospho-SEK1/MKK4 (Ser257) (C5) Rabbit mAb (PE Conjugate) SEK1MKK4S257-C5#2367, or concentration-matched Rabbit (G9) mAb IgG Isotype Control (PE Conjugate) #2142 for cells treated with lambda phosphatase (black) or treated with UV+TPA (blue).