


For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry	Anti-Rabbit IgG	Monoclonal	Rabbit IgGk

<b>Format:</b>	Unconjugated
<b>Cross Reactivity:</b>	Predicted to work with mouse, rat and other homologues.
<b>Formulation:</b>	1X PBS, 0.02% NaN3, 50% Glycerol, 0.1% BSA
<b>Preparation:</b>	Protein A+G
<b>Reactivity:</b>	Human,Mouse
<b>Recommended Usage:</b>	1μg/mL - 0.001μg/mL. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.
<b>Immunogen:</b>	A synthetic phospho-peptide corresponding to residues surrounding Ser2448 of human phospho mTOR
<b>Description:</b>	mTOR, <b>mammalian target of rapamycin</b> , is a Serine/Threonine protein kinase (1-2) that functions as an amino acid and ATP sensor to balance cell growth and nutrient availability (3-4). When sufficient nutrients are available, mTOR transmits a positive signal to p70 S6 kinase and participates in the inactivation of 4E-BP1 (5). mTOR plays a key role in homeostasis and cell growth, and phospho mTOR may be abnormally regulated in tumors. mTOR is a potential target for anti-cancer therapy (6).
<b>References:</b>	<ol style="list-style-type: none"><li>1. Sabers, C.J. et al. (1995) J. Biol. Chem. 270: 815-822.</li><li>2. Brown, E.J. et al. (1994) Nature. 369: 756-758.</li><li>3. Gingras, A.C. et al. (2001) Gene. Dev. 15: 807-826.</li><li>4. Dennis, P.B. et al. (2001) Science. 294: 1102-1105.</li><li>5. Fang, Y. et al. (2001) Science. 294: 1942-1945.</li><li>6. Huang, S. and Houghton, P.J. (2003) Curr. Op. Pharmacol. 3: 371-377.</li></ol>



Flow cytometric analysis of NIH3T3 cells secondary antibody only negative control (blue) or 0.01 µg/mL of isotype control Cat. #2141 (orange) or untreated (red) or treated with PDGF (green) using Phospho-mTOR (Ser2448) antibody mTORS2448-E11 at 0.01 µg/mL. Cat. #2376.