
Phospho-mTOR (Ser2448) (E11) rabbit mAb PE Conjugate

www.abwizbio.com Support: info@abwizbio.com Order: sales@abwizbio.com

Catalog: #2377 Store at: 2-8°C

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry	N/A	Monoclonal	Rabbit IgGk
Format:	PE		
Cross Reactivity:	Predicted to work with mouse, rat and other homologues.		
Formulation:	1X PBS, 0.09% NaN3, 0.2% BSA		
Preparation:	Protein A+G		
Reactivity:	Human,Mouse		
Recommended Usage:	For flow cytometric staining, the suggested use of this reagent is 5 μL per million cells or 5 μL per 100 μL of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.		
Immunogen:	A synthetic phospho-peptide corresponding to residues surrounding Ser2448 of human phospho mTOR		
Description:	mTOR, mammalian target of rapamycin, is a Serine/Threonine protein kinase (1-2) that functions as an amino acid and ATP sensor to balance cell growth and nutrient availability (3-4). When sufficient nutrients are available, mTOR transmits a positive signal to p70 S6 kinase and participates in the inactivation of 4E-BP1 (5). mTOR plays a key role in homeostasis and cell growth, and phospho mTOR may be abnormally regulated in tumors. mTOR is a potential target for anti-cancer therapy (6).		
References:	 Sabers, C.J. et al. (1995) J. Biol. Chem. 270: 815-822. Brown, E.J. et al. (1994) Nature. 369: 756-758. Gingras, A.C. et al. (2001) Gene. Dev. 15: 807-826. Dennis, P.B. et al. (2001) Science. 294: 1102-1105. Fang, Y. et al. (2001) Science. 294: 1942-1945. Huang, S. and Houghton, P.J. (2003) Curr. Op. Pharmacol. 3: 371-377. 		



Flow cytometric analysis of A431 cells treated with phosphatase and unstained as negative control (blue) or treated with phosphatase (red) or EGF (green) and stained using Phospho-mTOR (Ser2448) PE conjugated antibody mTORS2448-E11. Cat. #2377. © 2025 Abwiz Bio