

# Phospho-mTOR (Ser2448) (E11) rabbit mAb FITC Conjugate

[www.abwizbio.com](http://www.abwizbio.com)

Support: [info@abwizbio.com](mailto:info@abwizbio.com)

Order: [sales@abwizbio.com](mailto:sales@abwizbio.com)


**Catalog:** #2378

**Store at:** 2-8°C

*For Research Use Only. Not For Use In Diagnostic Procedures.*

Applications	Detection	Clonality	Isotype
Flow Cytometry	N/A	Monoclonal	Rabbit IgGk

<b>Format:</b>	FITC
<b>Cross Reactivity:</b>	Predicted to work with mouse, rat and other homologues.
<b>Formulation:</b>	1X PBS, 0.09% NaN <sub>3</sub> , 0.2% BSA
<b>Preparation:</b>	Protein A+G
<b>Reactivity:</b>	Human, Mouse
<b>Recommended Usage:</b>	For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells or 5 µL per 100 µL of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.
<b>Immunogen:</b>	A synthetic phospho-peptide corresponding to residues surrounding Ser2448 of human phospho mTOR
<b>Description:</b>	mTOR, mammalian target of rapamycin, is a Serine/Threonine protein kinase (1-2) that functions as an amino acid and ATP sensor to balance cell growth and nutrient availability (3-4). When sufficient nutrients are available, mTOR transmits a positive signal to p70 S6 kinase and participates in the inactivation of 4E-BP1 (5). mTOR plays a key role in homeostasis and cell growth, and phospho mTOR may be abnormally regulated in tumors. mTOR is a potential target for anti-cancer therapy (6).
<b>References:</b>	<ol style="list-style-type: none"><li>1. Sabers, C.J. et al. (1995) J. Biol. Chem. 270: 815-822.</li><li>2. Brown, E.J. et al. (1994) Nature. 369: 756-758.</li><li>3. Gingras, A.C. et al. (2001) Gene. Dev. 15: 807-826.</li><li>4. Dennis, P.B. et al. (2001) Science. 294: 1102-1105.</li><li>5. Fang, Y. et al. (2001) Science. 294: 1942-1945.</li><li>6. Huang, S. and Houghton, P.J. (2003) Curr. Op. Pharmacol. 3: 371-377.</li></ol>



Flow cytometric analysis of A431 cells treated with phosphatase and unstained as negative control (blue) or treated with phosphatase (red) or EGF (green) and stained using Phospho-mTOR (Ser2448) FITC conjugated antibody mTORS2448-E11. Cat. #2378.