

Phospho-mTOR (Ser2448) (E11) rabbit mAb APC Conjugate

www.abwizbio.com

Support: info@abwizbio.com

Order: sales@abwizbio.com

#2379

Store at: 2-8°C

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry	N/A	Monoclonal	Rabbit IgGk

Format: APC

Cross Reactivity: Predicted to work with mouse, rat and other homologues.

Formulation: 1X PBS, 0.09% NaN₃, 0.2% BSA

Preparation: Protein A+G

Reactivity: Human, Mouse

Recommended


Usage: For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells or 5 µL per 100 µL of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.

Immunogen: A synthetic phospho-peptide corresponding to residues surrounding Ser2448 of human phospho mTOR

Description: mTOR, mammalian target of rapamycin, is a Serine/Threonine protein kinase (1-2) that functions as an amino acid and ATP sensor to balance cell growth and nutrient availability (3-4). When sufficient nutrients are available, mTOR transmits a positive signal to p70 S6 kinase and participates in the inactivation of 4E-BP1 (5). mTOR plays a key role in homeostasis and cell growth, and phospho mTOR may be abnormally regulated in tumors. mTOR is a potential target for anti-cancer therapy (6).

References:

1. Sabers, C.J. et al. (1995) J. Biol. Chem. 270: 815-822.
2. Brown, E.J. et al. (1994) Nature. 369: 756-758.
3. Gingras, A.C. et al. (2001) Gene. Dev. 15: 807-826.
4. Dennis, P.B. et al. (2001) Science. 294: 1102-1105.
5. Fang, Y. et al. (2001) Science. 294: 1942-1945.
6. Huang, S. and Houghton, P.J. (2003) Curr. Op. Pharmacol. 3: 371-377.



Flow cytometric analysis of A431 cells treated with phosphatase and unstained as negative control (blue) or treated with phosphatase (red) or EGF (green) and stained using Phospho-mTOR (Ser2448) APC conjugated antibody mTORS2448-E11. Cat. #2379.