Phospho-NFKB p65 (Ser536) (B7) rabbit mAb

www.abwizbio.com
Support: info@abwizbio.com
Order: sales@abwizbio.com

Catalog: #2391

Store at: -20ºC

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry,WB	Anti-Rabbit IgG	Monoclonal	Rabbit IgGk

Format: Unconjugated

Cross Reactivity: Predicted to work with mouse, rat and other homologues.

Formulation: 1X PBS, 0.02% NaN3, 50% Glycerol, 0.1% BSA

Preparation: Protein A+G

Reactivity: Human


Recommended Usage: 1µg/mL – 0.001µg/mL. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.

Immunogen: A synthetic phospho-peptide corresponding to residues surrounding Ser536 of human phospho-NFKB p65

Description: The nuclear factor κB (NFκB)/Rel family of transcription factors play a pivotal role in inflammatory and immune responses (1,2). NF-kappa-B is present in almost all cell types and is involved in many biological processes including immunity, inflammation, cell growth and differentiation, apoptosis, and tumorigenesis. NFkB is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFkB1/p105, NFkB1/p50, REL and NFkB2/p52. The dimers bind at κB sites in the target gene DNA. Individual dimers have distinct preferences for different κB sites and can act as either transcriptional activators or repressors. NFkB Ser536 phosphorylation stimulates Lys310 acetylation and interaction of phospho NFkB with CBP. Acetylated/phospho NFkB induces enhanced transcriptional activity.


References:

1. Baeuerle, P.A. and Henkel, T. (1994) Annu. Rev. Immunol. 12:141-179.
2. Baeuerle, P.A. and Baltimore, D. (1996) Cell. 87:13-20.



©2026 Abwiz Bio

Flow cytometric analysis of HeLa cells unstained and untreated as negative control (blue) or stained and untreated (red) or stained and treated with TNFa plus CalA (green) using phospho-NFKB p65 (Ser536) antibody NFKBP65S536-B7 at 0.05 ug/mL Cat. #2391.



Western blot analysis of HeLa cell extract untreated or treated with TNFa plus CAlA using phospho-NFKB p65 (Ser536) antibody NFKBP65S536-B7 at 0.05 ug/mL. Cat. #2391.

©2026 Abwiz Bio