

# Phospho-HS1 (Tyr397) (F12) rabbit mAb

[www.abwizbio.com](http://www.abwizbio.com)

Support: [info@abwizbio.com](mailto:info@abwizbio.com)


Order: [sales@abwizbio.com](mailto:sales@abwizbio.com)

Catalog: #2396


Store at: -20°C

For Research Use Only. Not For Use In Diagnostic Procedures.


Applications	Detection	Clonality	Isotype
Flow Cytometry, WB	Anti-Rabbit IgG	Monoclonal	Rabbit IgGk
Format:	Unconjugated		
Cross Reactivity:	Predicted to work with mouse, rat and other homologues.		
Formulation:	1X PBS, 0.02% NaN3, 50% Glycerol, 0.1% BSA		
Preparation:	Protein A+G		
Reactivity:	Human, Mouse		
Recommended Usage:	1µg/mL – 0.001µg/mL. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.		
Immunogen:	A synthetic phospho-peptide corresponding to residues surrounding Tyr397 of human phospho HS1		
Description:	HS1 is expressed in lymphoid and hematopoietic cells, and is heavily post-translationally modified. HS1 deficient mouse models have demonstrated the protein's role in receptor-mediated apoptosis and proliferation. HS1 is phosphorylated at Tyr378 and Tyr397 by the kinase Syk, providing a high-affinity binding site for SH2 domains from the Src family. Following this interaction, HS1 is then phosphorylated at Tyr222 by c-Fgr, Lyn, and Fyn kinases. HS1 plays an important role in T cell signaling, where HS1 phosphorylation recruits and activates Vav1 at the immune synapse. As a homolog of the actin binding protein cortactin, HS1 has been shown to mediate neutrophil chemotaxis through phosphorylation of tyrosines 222, 378, and 397.		
References:	Brunati AM, Donella-Deana A, James P, Quadroni M, Contri A, Marin O, and Pinna LA. (1999) Journal of Biological Chemistry. 274:7557-7564. Cavnar PJ, Mogen K, Berthier E, Beebe DJ, and Huttenlocher A. (2012) Journal of Biological Chemistry. 287: 25466-25477.		



Flow cytometric analysis of Ramos cells secondary antibody only negative control (blue) or untreated (red) or treated with pervanadate (green) using Phospho-HS1 (Tyr397) antibody HS1Y397-F12 at 0.01 $\mu$ g/mL. Cat. #2396.




Western blot analysis of Ramos cell extract, untreated or treated with 300 nM Thapsigargin for 30 min using HS1 (Tyr397) antibody HS1Y397-F12 at 0.01  $\mu$ g/mL. Cat. #2396.




	IgG	Treatment	Peptide Block	Median : BL1-A
□	F12	Pv	Non-phos.	117301
□	F12	Ctrl	Non-phos.	863
□	F12	Pv	Phospho.	355
□	F12	Ctrl	Phospho.	268
□	F12	Pv	-	115459
□	F12	Ctrl	-	856
□	2' only	Ctrl	-	253

Peptide blocking flow cytometric analysis of Ramos cells secondary antibody only negative control (light blue) or untreated (red) or treated with pervandadate (green) or untreated and blocked with phospho-peptide (black) or treated and blocked with phospho peptide (gold) or untreated and blocked with non-phospho peptide (dark blue) or treated and blocked with non-phospho peptide (purple) using Phospho-HS1 (Tyr397) antibody HS1Y397-F12 at 0.01µg/mL. Cat. #2396.



Western blot analysis of Ramos cell extract untreated or treated with 300 nM thapsigargin for 30 min using 10 ng/mL Phospho-HS1 (Tyr397) antibody HS1Y397-F12 at 0.01µg/mL. Cat. #2396 or Company C antibody at 3 ng/mL (manufacturer's recommended concentration) developed using the same exposure.



Flow cytometric analysis of mouse cells secondary antibody only negative control (blue) or control (red) or stimulated (green) using Phospho-HS1 (Tyr397) antibody HS1Y397-F12 at 0.01µg/mL. Cat. #2396.