www.abwizbio.com Support: info@abwizbio.com Order: sales@abwizbio.com

Catalog: #2399

Store at: 2-8°C

For Research Use Only. Not For Use In Diagnostic Procedures.

| Applications Flow Cytometry | Detection N/A | Clonality Monoclonal | Isotype Rabbit IgGk |
|--------------------------------|---|--------------------------------|-------------------------------|
| | | Monocional | Rubble Igek |
| Format: | APC | | |
| Cross Reactivity: | Predicted to work with mouse, rat and other homologues. | | |
| Formulation: | 1X PBS, 0.09% NaN3, 0.2% BSA | | |
| Preparation: | Protein A+G | | |
| Reactivity: | Human,Mouse | | |
| Recommended Usage: | For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells or 5 μ L per 100 μ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information. | | |
| Immunogen: | A synthetic phospho-peptide corresponding to residues surrounding Tyr397 of human phospho HS1 | | |
| Description: | HS1 is expressed in lymphoid and hematopoietic cells, and is heavily post- translationally modified. HS1 deficient mouse models have demonstrated the protein's role in receptor-mediated apoptosis and proliferation. HS1 is phosphorylated at Tyr378 and Tyr397 by the kinase Syk, providing a high-affinity binding site for SH2 domains from the Src family. Following this interaction, HS1 is then phosphorylated at Tyr222 by c-Fgr, Lyn, and Fyn kinases. HS1 plays an important role in T cell signaling, where HS1 phosphorylation recruits and activates Vav1 at the immune synapse. As a homolog of the actin binding protein cortactin, HS1 has been shown to mediate neutrophil chemotaxis through phosphorylation of tyrosines 222, 378, and 397. | | |
| References: | Brunati AM, et al., (1999) Journal of Biological Chemistry. 274:7557-7564. Cavnar PJ, et al., (2012) Journal of Biological Chemistry. 287: 25466-25477. | | |

Phospho-HS1 (T397) APC

Flow cytometric analysis of Ramos cells untreated and unstained as negative control (blue) or untreated (red) or treated with pervanadate (green) and stained using Phospho-HS1 (Tyr397) APC conjugated antibody HS1Y397-F12. Cat. #2399.

