Phospho-Jak1 (Tyr1034/1035) (F11) rabbit mAb APC Conjugate

www.abwizbio.com

Support: info@abwizbio.com
Order: sales@abwizbio.com

#2414 Store at: 2-8°C

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Flow Cytometry	Anti-Rabbit IgG	Monoclonal	Rabbit IgGk

Format: APC

Cross Reactivity: Predicted to work with mouse, rat and other homologues.

Formulation: 1X PBS, 0.09% NaN3, 0.2% BSA

Preparation: Protein A+G

Reactivity: Human, Mouse

Recommended

Usage: For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells or 5 μ L per 100

µL of staining volume. It is recommended that the reagent be titrated for optimal performance for each

application. See product image legends for additional information.

Immunogen: A synthetic phospho-peptide corresponding to residues surrounding Tyr1034/1035 of human

phospho-Jak1

Description: <p>Jak1 plays an essential role in the IFN-? and IFN-? response pathways and is

tyrosine-phosphorylated upon cellular exposure to these signals. Jak1 oral inhibitors have been used to benefit patients with advanced myelofibrosis, where Jak1 was initially shown to be constitutively active in the peripheral blood cells of these patients. Targeted, small-molecule Jak inhibitors have also been used for treatment of rheumatoid arthritis. In cases of advanced melanoma, acquired resistance to PD-1 blockade drugs is associated with loss-of-functions of mutations in Jak1/2 genes. These mutations block interferon gamma signaling and prevent programmed death ligand 1 (PD-L1)

expression in tumor cells.</p>


References: Muller M, et al. (1993) Nature. 366: 129-135.

Verstovesk S, et al. (2010) New England Journal of Medicine. 363: 1117-1127.

Shin DS, et al. (2017) Cancer Discovery. 7:188-201.

Boyle DL, et al. (2014) Annals of the Rheumatic Diseases. 74: 1311-1316.





Phospho-JAK1(Y1034/1035) APC

Flow cytometric analysis of Jurkat cells untreated (red) or treated with IFN? + IL-4 + pervanadate (green) using Phospho-Jak1 (Tyr1034/1035) (F11) Rabbit mAb (APC Conjugate) Jak1Y10341035-F11 #2414, or concentration-matched Rabbit (G9) mAb IgG Isotype Control (APC Conjugate) #2144 for cells untreated (black) or treated with IFN? + IL-4 + pervanadate (blue).