Phospho-Lamin A/C (Ser22) (CF12) rabbit mAb PE conjugate

www.abwizbio.com

Support: info@abwizbio.com Order: sales@abwizbio.com


#2437

Store at: 2-8°C

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	s Detection	Clonality	Isotype
Flow Cytomet	ry N/A	Monoclonal	Rabbit IgGk
Format:	PE		
Cross Reactivity:	Predicted to work with mouse, rat and other homologues.		
Formulation:	1X PBS, 0.09% NaN3, 0.2% BSA		
Preparation:	Protein A+G		
Reactivity:	Human,Mouse		
Recommended Usage:	For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells or 5 μ L per 100 μ L of staining volume. It is recommended that the reagent be titrated for optimal performance for each application. See product image legends for additional information.		
Immunogen:	A synthetic phospho-peptide corresponding to residues surrounding Ser22 of human phospho Lamin A/C		
Description:	Lamins are nuclear membrane proteins that are involved with cell cycle control, chromatin organization, and DNA replication (1,2). Upon caspase 6 cleavage, lamin A/C serves as a molecular marker for capspase 6 activity. Lamin A/C is cleaved during cellular apoptosis into a large (41-50 kDa) as well as a small (28 kDa) fragment leading to nuclear dysfunction and cell death (3) Phosphorylation of lamin A/C at Ser32 by MAPK/CDK signals cell cycle progression and mitosis (4,5).		
References:	 Goldberg, M, et al., (1999) Crit Rev Eu Yabuki, M., et ql., (1999) Physiol Cher Rao, L et al., (1996) J Cell Biol 135: 14 Orth K, et al., (1996) J Biol Chem 271: Nousiainen M, et al., (2006) Proc Natl 	n Phys Med NMR 31: 77-84 441:55. 16443-6	





Flow cytometric analysis of Hela cells untreated (red) or treated with nocadozole (green) using Phospho-Lamin A/C (Ser22) (CF12) Rabbit mAb (PE Conjugate) LaminACS22-CF12 #2437, or concentration-matched Rabbit (G9) mAb IgG Isotype Control (PE Conjugate) #2142 for cells untreated (black) or treated with nocodazole (blue).

Phospho-Lamin A/C (S22) PE

