



**Catalog:** #2491**Store at:** 2-8°C*For Research Use Only. Not For Use In Diagnostic Procedures.*

Applications	Detection	Clonality	Isotype
ELISA	Anti-Rabbit IgG	Monoclonal	Rabbit IgGk

<b>Format:</b>	Unconjugated
<b>Cross Reactivity:</b>	Highly-specific to SARS-CoV-2 nucleoprotein; does not cross-react with nucleoprotein from SARS-CoV or other coronaviruses.
<b>Formulation:</b>	1X PBS, 0.02% NaN3
<b>Preparation:</b>	Protein A
<b>Reactivity:</b>	Other
<b>Recommended Usage:</b>	Used for specific, high-sensitivity detection of SARS-CoV-2 nucleocapsid protein (NP) in immunoassay. Can be paired with other NP-specific clones for detection in sandwich ELISA format.
<b>Immunogen:</b>	SARS-CoV-2 nucleoprotein (NP) specific peptide
<b>Description:</b>	We have leveraged our next-generation rabbit mAb discovery platform to develop recombinant rabbit monoclonal antibodies with extremely high sensitivity and specificity to SARS-CoV-2. These antibodies, which include clones 75G5a ( <a href="#">Abwiz Cat. #2481</a> ), 84C4a ( <a href="#">Abwiz Cat. #2486</a> ), 84D7 (Abwiz Cat. #2491), and 85C1 ( <a href="#">Abwiz Cat. #2496</a> ) can be paired in sandwich detection assay and used to detect nucleoprotein (NP) antigen from SARS-CoV-2. Antibodies 75G5a, 84C4a, and 85C1 do not cross-react to the highly related SARS-CoV virus or to any other coronaviruses tested. Sandwich ELISA detection using TMB/acid developer reliably detects NP antigen in the pg/mL range, and sensitivity is expected to be even higher when using more sensitive developer strategies. This panel of antibodies can be used as raw materials for diagnostic kits and can be applied to lateral flow systems for diagnostic detection of COVID-19.

**References:**





Microtiter wells were coated with SARS-CoV-2 (COVID-19) Nucleocapsid Protein (NP) and SARS-CoV NP at 1  $\mu$ g/mL. Purified rabbit monoclonal antibody 84D7 (Cat# 2491) was serially diluted 1:2 starting at 1  $\mu$ g/mL, and shows very strong and specific binding to COVID-19 NP antigen, with no significant cross-reactivity to SARS-CoV NP antigen.

Clone	$K_D$ (M)	$K_{on}$ (1/Ms)	$K_{off}$ (1/s)
75Ga	$8.25 \times 10^{-11}$	$1.36 \times 10^5$	$1.12 \times 10^{-5}$
84C4a	$9.38 \times 10^{-11}$	$1.53 \times 10^5$	$1.43 \times 10^{-5}$
84D7	$2.23 \times 10^{-10}$	$7.43 \times 10^4$	$1.66 \times 10^{-5}$

Affinity measurement of SARS-CoV-2 antibody (84D7)