

SARS-CoV2 LxC1-G4 human neutralizing mAb

www.abwizbio.com

Support: info@abwizbio.com

Order: sales@abwizbio.com

#2546

Store at: -20°C

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications	Detection	Clonality	Isotype
Functional Assay,ELISA	Anti-SARS-CoV-2 NP mAb	Monoclonal	Rabbit IgGk

Format: Unconjugated

Cross Reactivity: No

Formulation: 1X PBS

Preparation: Protein A


Reactivity: Other

Recommended Usage: Experimental conditions may be optimized.


Immunogen: A recombinant SARS-CoV-2 spike protein containing RBD.

Description: N/A


References: N/A




Spike protein trimers were coated at 2 µg/mL and incubated with serially diluted antibodies. The bound antibodies were detected with Peroxidase AffiniPure Goat Anti-Human IgG, Fc γ fragment specific (Jackson Immuno Research Catalog#109-035-098) or Peroxida




Spike protein trimers were coated at 2 µg/mL and incubated with serially diluted antibodies. The bound antibodies were detected with Peroxidase AffiniPure Goat Anti-Human IgG, Fc? fragment specific (Jackson Immuno Research Catalog#109-035-098) or Peroxida




Spike protein trimers were coated at 2 µg/mL and incubated with serially diluted antibodies. The bound antibodies were detected with Peroxidase AffiniPure Goat Anti-Human IgG, Fc? fragment specific (Jackson Immuno Research Catalog#109-035-098) or Peroxida




Spike protein trimers were coated at 2 µg/mL and incubated with serially diluted antibodies. The bound antibodies were detected with Peroxidase AffiniPure Goat Anti-Human IgG, Fc? fragment specific (Jackson Immuno Research Catalog#109-035-098) or Peroxida




Spike protein trimers were coated at 2 µg/mL and incubated with serially diluted antibodies. The bound antibodies were detected with Peroxidase AffiniPure Goat Anti-Human IgG, Fc? fragment specific (Jackson Immuno Research Catalog#109-035-098) or Peroxida



Spike protein trimers were coated at 2 µg/mL and incubated with serially diluted antibodies. The bound antibodies were detected with Peroxidase AffiniPure Goat Anti-Human IgG, Fc? fragment specific (Jackson Immuno Research Catalog#109-035-098) or Peroxida



Spike protein trimers were coated at 2 µg/mL and incubated with serially diluted antibodies. The bound antibodies were detected with Peroxidase AffiniPure Goat Anti-Human IgG, Fc? fragment specific (Jackson Immuno Research Catalog#109-035-098) or Peroxida



Spike protein trimers were coated at 2 µg/mL and incubated with serially diluted antibodies. ACE2-Avitag protein (Acro Biosystems AC2-H82E6) was added to 0.037 µg/mL and the bound ACE2 was detected with High Sensitivity StreptAvidin-HRP (Thermo Fisher Sci

IC50

N/A

	Wild type (WT)	Alpha (B.1.1.7)	Beta (B.1.351)	Gamma (P.1)	Kappa (B.1.617.1)	Epsilon (B.1.427)	Epsilon (B.1.429)
LxC1-G4	22.38072	69.60794	38.41653	20.40104	13.08920	19.40038	12.99241